

N-of-1 trial of serial induced cough challenges in COVID-19 showing an interplay between Nrf2, TRPA1 and TRPV1 agonists

Jean Bousquet^{1,2}, Wienczyslawa Czarlewski^{3,4}, Torsten Zuberbier¹, Joaquim Mullol⁵, Hubert Blain⁶, Jean-Paul Cristol⁷, Rafael de la Torre⁸⁻¹⁰, Nieves Pizarro Lozano⁹, Vincent Le Moing¹¹, Anna Bedbrook^{2,4}, Ioana Agache¹², Cezmi A Akdis¹³, G Walter Canonica¹⁴, Alvaro A Cruz¹⁵, Alessandro Fiocchi¹⁶, Joao Fonseca¹⁷, Susana Fonseca¹⁸, Bilun Gemicicoglu¹⁹, Tari Haahtela²⁰, Guido Iaccarino²¹, Juan-Carlos Ivancevich²², Marek Jutel²³, Ludger Klimek²⁴, Helga Kraxner²⁵, Piotr Kuna²⁶, Désirée Larenas-Linnemann²⁷, Erik Melén²⁸, Yoshitaka Okamoto²⁹, Nikolaos G Papadopoulos³⁰, Oliver Pfaar³¹, Frederico Regateiro³², Jacques Reynes¹¹, Yves Rolland³³, Philip P Rouadi³⁴, Boleslaw Samolinski³⁵, Sanna Toppila-Salmi²⁰, Arunas Valiulis³⁶, Hak-Jong Choi³⁷, Hyun Ju Kim³⁸, Josep M Anto^{9,10,39,40}

1. Charité, Universitätsmedizin Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Comprehensive Allergy Center, Department of Dermatology and Allergy, Berlin, Germany
2. University hospital and MACVIA France, Montpellier, France
3. Medical Consulting Czarlewski, Levallois, France.
4. MASK-air, Montpellier, France.
5. Rhinology Unit & Smell Clinic, ENT Department, Hospital Clinic - Clinical & Experimental Respiratory Immunology, IDIBAPS, CIBERES, Universitat de Barcelona, Barcelona, Spain
6. Department of Geriatrics, Montpellier University Hospital, Montpellier, France
7. Laboratoire de Biochimie et Hormonologie, PhyMedExp, Université de Montpellier, INSERM, CNRS, CHU de Montpellier, France
8. CIBER Fisiopatología de la Obesidad y Nutrición (CIBEROBN), Madrid, Spain
9. IMIM (Hospital del Mar Research Institute), Barcelona, Spain
10. Universitat Pompeu Fabra (UPF), Barcelona, Spain
11. Maladies Infectieuses et Tropicales, CHU Montpellier, France
12. Faculty of Medicine, Transylvania University, Brasov, Romania
13. Swiss Institute of Allergy and Asthma Research (SIAF), University of Zurich - Christine Kühne - Center for Allergy Research and Education (CK-CARE), Davos, Switzerland
14. Personalized Medicine, Asthma and Allergy, Humanitas Clinical and Research Center IRCCS and Department of Biomedical Sciences, Humanitas University, Pieve Emanuele (MI), Italy
15. Fundação ProAR, Federal University of Bahia and GARD/WHO Planning Group, Salvador, Brazil
16. Division of Allergy, Department of Pediatric Medicine - The Bambino Gesù Children's Research Hospital Holy see, Rome, Italy
17. CINTESIS, Center for Research in Health Technologies and Information Systems, Faculdade de Medicina da Universidade do Porto, Porto, Portugal and MEDIDA, Lda, Porto, Portugal
18. GreenUPorto - Sustainable Agrifood Production Research Centre, DGAOT, Faculty of Sciences, University of Porto, Campus de Vairão, Vila do Conde, Portugal
19. Department of Pulmonary Diseases, Istanbul University-Cerrahpasa, Cerrahpasa Faculty of Medicine, Istanbul
20. Skin and Allergy Hospital, Helsinki University Hospital, and University of Helsinki, Helsinki, Finland
21. Department of Advanced Biomedical Sciences, Federico II University, Napoli, Italy
22. Servicio de Alergia e Inmunología, Clínica Santa Isabel, Buenos Aires, Argentina
23. Department of Clinical Immunology, Wrocław Medical University and ALL-MED Medical Research Institute, Wrocław, Poland
24. Center for Rhinology and Allergology, Wiesbaden, Germany
25. Department of Otorhinolaryngology, Head and Neck Surgery, Semmelweis University, Budapest, Hungary
26. Division of Internal Medicine, Asthma and Allergy, Barlicki University Hospital, Medical University of Lodz, Poland
27. Center of Excellence in Asthma and Allergy, Médica Sur Clinical Foundation and Hospital, México City, Mexico
28. Institute of Environmental Medicine, Karolinska Institutet and Sachs' Children's Hospital, Stockholm, Sweden
29. Dept of Otorhinolaryngology, Chiba University Hospital, Chiba, Japan
30. Division of Infection, Immunity & Respiratory Medicine, Royal Manchester Children's Hospital, University of Manchester, Manchester, UK, and Allergy Department, 2nd Pediatric Clinic, Athens General Children's Hospital "P&A Kyriakou," University of Athens, Athens, Greece
31. Department of Otorhinolaryngology, Head and Neck Surgery, Section of Rhinology and Allergy, University Hospital Marburg, Philipps-Universität Marburg, Germany
32. Allergy and Clinical Immunology Unit, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal, Faculty of Medicine, Institute of Immunology, University of Coimbra, Coimbra, Portugal. Faculty of Medicine, ICBR - Coimbra Institute for Clinical and Biomedical Research, CIBB, University of Coimbra, Coimbra, Portugal.

33. CHU Toulouse, France
34. Department of Otolaryngology-Head and Neck Surgery, Eye and Ear University Hospital, Beirut, Lebanon
35. Department of Prevention of Environmental Hazards and Allergology, Medical University of Warsaw, Poland
36. Vilnius University Faculty of Medicine, Institute of Clinical Medicine & Institute of Health Sciences, Vilnius, Lithuania
37. Microbiology and Functionality Research Group, Research and Development Division, World Institute of Kimchi, Gwangju, Korea
38. SME Service Department, Strategy and Planning Division, World Institute of Kimchi, Gwangju, Korea
39. CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain
40. ISGlobAL, Barcelona, Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Spain

Short Title: Nutrients acting on cough challenges in COVID-19

Abstract

Background: COVID-19 is commonly associated with impaired redox homeostasis responsible for reactive oxygen species accumulation. Antioxidants have been proposed as being effective in controlling COVID-19 symptoms. Nuclear factor (erythroid-derived 2)-like 2 (Nrf2) is the most effective antioxidant mechanism. TRPA1 (transient receptor potential ankyrin 1) is highly sensitive to oxidative stress and induces several COVID-19 symptoms. TRPA1 and TRPV1 (transient receptor potential vanillin 1) are potential candidates for COVID-19 symptoms.

Methods: Forty-nine open label induced cough challenges were carried out in an n-of-1 trial. Nutrients with various Nrf2 and TRPA1/TRPV1 agonist activity were used: broccoli seeds (potent Nrf2 agonist and mild TRPA1 agonist), berberine (Nrf2 only), black pepper, curcumin, ginger, green tea, resveratrol, zinc (potent TRPA1 activity and variable Nrf2 agonist) and red pepper (potent TRPV1 agonist).

Results: Berberine and zinc were not effective. All other nutrients except resveratrol were rapidly effective (1-10 minutes). The effect of green tea, curcumin + black pepper, ginger, resveratrol or red pepper disappeared within 1 to 4 hours. Broccoli had a longer duration of action (5-7 hours). The duration of the effect increased to around 10 hours when low doses of curcumin + black pepper or resveratrol were added to broccoli. Paracetamol low dose (its metabolite N-acetyl-p-benzoquinone imine is a TRPA1/TRPV1 agonist) increases the duration of action of combinations to over 14 hours.

Conclusions: A synergy between Nrf2 antioxidative effect and TRPA1 desensitization (and TRPV1) is proposed. Research should confirm this trial and find the mechanisms of synergy.

Key words: COVID-19, Nrf2, TRPA1, TRPV1, channel desensitization, sulforaphane, cough

Introduction

Three stages of COVID-19 have been described: (i) a viral infection lasting for one to two weeks; (ii) a second phase characterized by an intertwined cytokine and oxidative stress storm, independent of infection; and (iii) a recovery phase that may last for some months. Many patients during the recovery phase suffer from persistent symptoms including cough.

A common denominator in all conditions associated with COVID-19 appears to be the impaired redox homeostasis responsible for reactive oxygen species (ROS) accumulation ¹. Many foods have antioxidant properties and many mechanisms may be involved. However, the activation of the nuclear factor (erythroid-derived 2)-like 2 (Nrf2) the most potent antioxidant transcription factor may be of primary importance.² Differences in COVID-19 death rates among countries may in part associated with Nrf2 and Nrf2-interacting nutrients like fermented vegetables could reduce COVID-19 severity ³⁻⁵. Many Nrf2-interacting nutrients such as spices are TRPA1 (transient receptor potential ankyrin 1) and/or TRPV1 (transient receptor potential vanillin 1) agonists ⁶. TRPA1, an excitatory ion channel, plays a pivotal role in augmenting sensory or vagal nerve discharges, evoking several COVID-19 symptoms ⁶. It is highly sensitive to oxidative stress. In COVID-19, rapid desensitization of TRAP1 and/or TRPV1 may reduce symptom severity ^{7 8}.

The gold-standard study design to assess intervention effectiveness is the population-based, parallel-group, randomized-controlled trial. However, n-of-1 trials in single patients are increasingly being relevant as a decision support tool to inform individualized treatment selection ⁹. The Oxford Center for Evidence-Based-Medicine ranks n-of-1 trials as level 1 evidence to determine if a treatment benefits a patient ¹⁰. These trials allow the patient to be involved in the selection of treatments. In an n-of-1 trial, the single patient alternate between treatments, not necessarily in a double-blind fashion ¹¹. N-of-1 trials are more useful in patients with stable conditions, with rapid onset effect of the treatment and a rapid washout ¹². N-of-1 trials are suited for nutritional interventions ¹³ and the hypothesis raised may represent a proof-of-concept that will be further confirmed by RCTs. Following the three clinical cases on COVID-19 treatment with nutrients ¹⁴, it appeared that N-of-1 trials were particularly suited to develop hypotheses for nutrients in the control of COVID-19 symptoms.

The present report consists of the self-description of an n-of-1 trial in COVID-19 case where the patient – who is also the author - describes induced cough challenges carried out during the recovery phase. The patient had moderate COVID symptoms during the first and second phases of COVID-19 ¹⁴. The patient presented persisting cough and nasal obstruction up to 65 days after the onset of COVID-19 symptoms. The challenges were carried out from 10 to 50 days after the cytokine storm with nutrients interacting with Nrf2, TRPA1 and TRPV1 aiming at differentiating time course of the effects and putative mechanisms.

1. Methods

1.1. Compounds tested

Broccoli was chosen as it contains glucoraphanin, a precursor of sulforaphane the most potent Nrf2 natural activator ^{15,16}. Broccoli seed capsules were labelled as containing 10% glucoraphanin (broccoli) but, using 3 different extracting methods ^{17,18} and mass spectrometry, the glucoraphanin content was 3% (around 7 μ mol per 100 mg of broccoli). Broccoli contains, selenium and the polyphenols such as kaempferol, quercetin glucosides and isorhamnetin, Nrf2 and TRPA1 agonists ¹⁹. These capsules were

shown to be effective in 3 patients during the first two phases of COVID-19¹⁴. However, it is difficult to deliver sulforaphane in an enriched and stable form for human consumption²⁰. Thus, glucoraphanin, the precursor of sulforaphane, is administered orally with myrosinase, the enzyme that transforms glucoraphanin into sulforaphane.

All nutrients tested have a variable Nrf2 agonist activity. Although the specificity of the different agonists is not clearcut it can be considered that curcuma and black pepper²¹, ginger, green tea²², quercetin and the plant polyphenol resveratrol²³ are all TRPA1 agonists, whereas berberine does not appear to have such an effect. Capsaicin from red pepper is the agonist of TRPV1 but it also has some TRPA1 activity. Zinc acts on Nrf2²⁴ and TRPA1²⁵.

Paracetamol increased the duration of efficacy (but not the efficacy) of broccoli treatment in three clinical cases¹⁴. The electrophilic metabolites N-acetyl-p-benzoquinone imine (NAPQI, hepatotoxic metabolite) and p-benzoquinone, but not paracetamol itself, activate TRPA1²⁶ and TRPV1²⁷. NAPQI also directly activates Nrf2²⁸, and benzoquinone desensitizes TRPA1²⁹. The physiological and toxicological responses of paracetamol form a continuum coordinated by the Wnt and Nrf2 pathways. Therapeutic doses produce reactive ROS and NAPQI in the cytoplasm but result in little permanent damage³⁰.

The compounds used in the study are listed in Table 1.

Table 1: Compounds used in the study

Nutrient	Origin	Composition as indicated on the label and recommended daily dose	N challenges	Dose
Broccoli	Pileje, Paris, France	Broccoli seeds 300 mg Glucoraphanin 3 % (measured)	9+11	150 mg 300 mg
Ginger	Arkopharma, Carros, France	Gingerol 6.3 mg	1+2	6.3 mg 12.6 mg
Green tea (loose or bag)	Lipton, Antwerp, Belgium		3	3.0 g 4.5 g
Green tea capsules	Arkopharma, Carros, France	1750 mg green tea	1+2	1750 mg
Curcumin + black pepper		Curcuma powder 560 mg Curcumin 103 mg Black pepper 16 mg Piperine 15 mg	2+4	103+16 mg 51.5+7.5 mg
Quercetin	Dynaveo, St Gilles, France	500 mg, extracted from <i>Sophora japonica</i> , (HPLC tested)	1	500 mg
Resveratrol		600 mg, extracted from <i>Polygonum cuspidum</i> (HPLC tested)	3+5	600 mg 1200 mg
Berberine		1200 mg, extracted from <i>Berbera aristata</i> (HPLC tested)	2	1200 amg 2400 mg
Zinc	Rubozinc®, Lacbatal, France	Zinc gluconate, 15 mg of Zn	1+1	15 mg
Paracetamol	Sanofi	Up to 3 g per day	1+4*	250 mg 500 mg

N: administered alone or (+) in combination, * some compounds were given in combination

1.2. Induced cough challenges

1.2.1. Characteristics of the patient

The patient is allergic to grass pollen and cat, and has moderate to severe allergic rhinitis which is perfectly controlled by as needed medications. Further to the onset of COVID-19, he experienced mild to moderate nasal obstruction (VAS up to 6/10) and proposed a score for his symptoms according to the VAS for allergic rhinitis (from 0 to 10) ^{31,32}. The patient also suffers from mild intermittent asthma, especially when exposed to high levels of air pollution. He uses formoterol-budesonide low dose when needed and his maximal annual consumption is around 40 doses. He has a severe bronchial hyperreactivity. He used formoterol-budesonide low dose to reduce COVID-19 cough without benefit.

1.2.2. Design of the test

Many scores exist for cough ³³. A visual analogue scale (VAS) score for “spontaneous cough” was used. When the patient had tracheobronchial pruritus, a deep inspiration followed by a rapid expiration led to immediate “wheezing” (not similar to asthma but similar to exposure to high levels of air pollution) and cough that were scored. He used a score for induced cough during challenge. For spontaneous cough during challenge, he decided to count the number of coughs per episode (Table 2).

Table 2: Cough symptom score during challenge ¹⁴

Spontaneous cough	Score	Induced cough after fast and hard expiration	Score
≥5 coughs for one episode	3	persisting severe cough	9-10
2-4 coughs for one episode	2	>9 coughs or intolerable	8
1 cough for one episode	1	6-9 coughs	7
		3-6 coughs, mild	6
		2-3 coughs, severe	5
		2-3 coughs, mild	4
		1 cough	3
		Loud “wheezing”	2
		Some “wheezing”	1

1.2.3. Open labelled induced cough challenges

In order to test the induced cough challenge and the broccoli dose that could be effective, the patient performed three induced cough challenges with a dose of 300 mg of broccoli (daily recommended dose). Similar to the FEV₁, he took the deepest breath he could, exhaled as fast and as hard as possible and counted the number of coughs according to Table 2. He found that induced cough improved within 10 minutes after broccoli ingestion and nasal obstruction even more rapidly. He then selected this dose for the double-blind, placebo-controlled (DB-PC) challenges (reported in a paper submitted and available online ¹⁴) and 10 pre-study open challenges. The DB-PC challenges (N=4 with broccoli and N=4 with lactose) and the open challenges showed similar data (Figure 1).

After more 60 days since the cytokine storm, the patient is still suffering from persistent cough and nasal obstruction, and needs the broccoli or other nutrient capsules when VAS for spontaneous cough >7/10. These episodes occur at around twice a day under treatment.

The patient performed a total of nine challenges carried out with broccoli capsules and 31 with Nrf2 and/or TRPA1-TRPV1 interacting nutrients (Table 1). Broccoli challenges were used to assess the

reproducibility of the challenges and to test whether the tracheobronchial reactivity was similar over time since improvement of the disease was not predictable.


The study was not double-blind since (i) the doses of some nutrients were up to 4 times higher than those recommended and there was a need to assess safety, and (ii) depending on the results obtained with the compound, the patient was able to design further challenges. In the induced cough challenges, symptoms were measured 5 minutes before dosing and at 1, 2, 5, 8, 15, 20, 25, 30, 45 and 60 minutes after. In many cases, the test was continued until induced cough scores were ≥ 7 , and induced cough challenges were repeated every hour. It was *a priori* proposed that the cutoff score level was 3/10 to discriminate a positive and a negative test. Induced cough, spontaneous cough and nasal obstruction were recorded throughout the study.

2. Results

2.1. Broccoli challenges

The results of the 9 broccoli challenges were very similar to both the 4 DB-PC broccoli challenges and the 10 open challenges performed before the onset of the study. This indicated that (i) there was a high reproducibility of the challenges and (ii) the patient had retained the same reactivity to induced cough throughout the study (Figure 1). Within 5-10 minutes after dosing, induced cough scores decreased from 7-8 to 1-3 and this was sustained for the following 50 minutes. A lower broccoli dose elicited the same effect.

Figure 1: Induced cough challenges with broccoli seed capsules



2.2. Rapid onset efficacy of Nrf2/TRAP1/TRPV1 nutrients

Several nutrients and paracetamol were used in the study (Table 2). Induced cough score was severe (score $\geq 7/10$) in all challenges before dosing. There was no effect for low-dose green tea (N=2), high dose berberine (N=2) or regular dose zinc (N=2). A significant effect (score ≤ 3) was observed within 2


minutes for curcumin + black pepper (N=2), ginseng (N=2); red pepper (N=3); 8-10 minutes for broccoli (N=9) and high-dose green tea (N=2), 20 min for paracetamol (N=1) and 30-45 minutes for resveratrol (N=3) (Figure 2 and Figure 3).

When low-dose broccoli (150 mg) was combined with low or regular doses of green tea capsules (N=2), half dose of ginger (N=2), half doses of curcumin + black pepper (N=2), or half doses of red pepper (N=2), the effect on cough began one minute after dosing and VAS was below 3 in 2 minutes. When combined with low-dose resveratrol, the effect on cough began at 2 minutes (N=5) (Figure 2,).

For nasal symptoms assessed before the cough challenges (Figure 3), the effect (nasal VAS ≤ 2) was observed within 1 minute for curcumin + pepper and higher doses of green tea, 2-5 minutes for broccoli, 25 min for paracetamol and 30-45 minutes for resveratrol. There was little effect for lower doses of green tea or zinc and no effect for berberine. All combinations - except those with green tea capsules - were rapidly effective. The two challenges with broccoli + green tea capsules were only followed by a small reduction in nasal symptoms (VAS 3) for the first 45 minutes.

Figure 2: Open labelled induced cough challenges with Nrf2-TRPA1/V1 interacting nutrients

Open label induced cough challenges with Nrf2-TRAP1/V1 nutrients



The arrows indicate the increased duration of adding broccoli to curcumin/black pepper (blue) and adding paracetamol to broccoli and curcumin/black pepper (grey).

2.3. Impact of Nrf2/TRAP1/TRPV1 nutrients during the first hour after dosing

The induced cough total score for each compound was compared with the mean values of the pre-study challenges carried out with 10 broccoli capsules (300 mg) (Figure 3). The 9 broccoli challenges showed similar results for 300 and 150 mg. On the other hand, the scores were higher for green tea but, more importantly, for berberine, paracetamol, resveratrol and zinc. Conversely, the scores were far lower for curcumin + black pepper, ginger, red pepper and all combinations including those with resveratrol.

Results differed for spontaneous cough occurring between challenges. Broccoli capsules had the lowest scores followed by curcumin + black pepper, most combinations, green tea, berberine and resveratrol.

Figure 3: Global results

	Product	Dose	Active metabolite (dose)	Mechanism	Induced cough challenge				Side effects (VAS)
					Induced cough 0-60	Spont. cough 0-60	Nasal VAS 0-60	Duration (min)	
	Broccoli (N=10)*	300 mg	Glucoraphanine 30 mg Kaemferol	Nrf2>>TRP ³⁴⁻³⁶	32				
1	Broccoli				38	4	34		
2	Broccoli				38	4	35		
3	Broccoli				30	5	25		
4	Broccoli				27	5	26	420	
5	Broccoli				32	4	28		
6	Broccoli				35	8	31	300	
7	Broccoli				36	7	28		
8	Broccoli	150mg	Glucoraphanine 15 mg		36	4	42	360	
9	Broccoli				32	4	45		
10	Berberine	1.2 g		Nrf2>TRP ^{37,38}	63	33	29	60	
11	Berberine	4.8 g			76	43	45	60	
12	Curcuma + black pepper	103 mg 16 mg	Curcumin Piperine	Nrf2-TRPA1 curcuma ^{21,34,39,40} peper ⁴¹⁻⁴³	17	4	15	240	HB: 2
13	Curcuma + black pepper	51 mg 8 mg	Curcumin Piperine		26	10	32	90	HB: 2
14	Ginger 2 caps	300 mg	Gingerol 12.6 mg	Nrf2-TRP ⁴⁴⁻⁴⁶	15	5	13	240	
15	Green tea bag	3 g		Nrf2>TRP ^{22,47}	54	12	34		
16		4.5 g			35	13	31		
17	Green tea capsule	1.75 g			47	17	27	60	
18	Green tea loose	unknown			54	12	33		
19	Quercetin	2 g		Nrf2>TRP ^{34,36,48}	78	21	53	60	
20	Red pepper	10 mg	Capsaicin	Nrf2<<TRPV1 ^{41,49} 42,43	21	11	18	120	HB: 2
21		20 mg			17	4	22	180	HB: 3
		30 mg			14	4	19	300	HB: 4
23	Resveratrol	600 mg		Nrf2>TRP ^{34,38,50}	63	14	46		
24		600 mg			68	32	38	300	
25		1200 mg			68	42	44	360	
26	Zinc	15 mg		Nrf2=TRP ^{24,25,51}	67	34	46	60	HB: 2
26	Paracetamol metabolites	500 mg		TRPA1-TRPV1 ^{26,29}	53	34	37	180	HB: 2
27	BG 150 + CU 50 + BP 8				13	6	14	500	HB: 2
28	BG 150 + CU 50 + BP 8				10	4	15	440	HB: 3
29	BG 150 + Ginger 150				19	3	14		
30	BG 150 + Res 300				20	7	19	600	
31	BG 300 + Res 300				20	11	22	600	
32	BG 300 + Res 300				16	12	26		
33	BG 150 + GTC 1700				22	3	24		
34	BG 150 + GTC 3250				8	14	29	600	
35	BG 150 + CU 50 + BP 8 + RP 20				5	6	11	780	HB: 4
36	BG 150 + CU 50 + BP 8 + paracetamol 250				6	7	18	900	HB: 2
37	BG 150 + CU 50 + BP 8 + paracetamol 250				11	2	16	720	
38	BG 150 + RP 20 + paracetamol 250				5	4	12	1020	HB: 6 GIP: 2*
39	BG 150 + Ginger 150 + paracetamol 250				11	2	13	720	HB: 3

BG: Broccoli, CU: curcumin, BP: Black pepper, Res: Resveratrol, GTC: Green tea capsule, RP: Red pepper

From 27 to 39: doses are in mg

Induced cough score (0-10*)	<20	20 to 29	30 to 39	40 to 59	≥59
Spontaneous cough score (0-10*)	<5	5 to 9	10 to 19	20 to 39	≥40
Nasal VAS (0-10*)	<15	15 to 19	20 to 29	30 to 39	≥40
Duration (min)	>660	421 to 660	181 to 420	60 to 180	≤60
*: 0: best, 10: worst					

Nasal symptoms were assessed before the cough challenge. Nasal total score (VAS) was similar for broccoli 300 mg, green tea and low-dose curcumin + pepper. It was lower for curcumin + pepper high dose and for all combinations. It was higher for broccoli 150 mg, berberine, resveratrol, zinc or paracetamol. A dose-response relationship was found for curcumin + black pepper or red pepper in nasal symptoms. There was a dose-dependent effect for broccoli (150 and 300 mg), curcumin + black pepper and red pepper.

2.4. Duration of the benefits of Nrf2/TRAP1 nutrients

In 27 cough challenges, a follow up was carried out until the patient experienced spontaneous cough (VAS≥7) or a cough score ≥7 during the challenge (Figure 3). The duration of efficacy was ranging from 300 to 420 minutes in the broccoli challenges, whatever the dose ingested. It was lower for green tea, resveratrol and particularly in curcumin + pepper, red pepper or paracetamol with a dose-dependent effect. For all broccoli combinations without paracetamol, the duration of action ranged from 440 to 600 minutes except for the combination with red pepper (780 minutes). For all broccoli combinations with paracetamol the duration of action ranged from 720 to 1020 minutes.

2-5- Symptoms experienced after the first hour

Representative examples of symptoms experienced after the first hour are presented in Figure 2. Induced cough was relatively well controlled (score ranging from 0 to 2 or 3/10) until the patient had a severe score with rapid onset. Nasal symptoms were similar to induced cough and spontaneously increased when the patient lost control of the disease.

During all of these challenges, there were some rapid, transient (less than one to 3 minutes) and mild episodes of tracheal irritation followed by one or two cough episodes and some nasal obstruction. These episodes were relatively common within the first hour following the ingestion of the compounds. At the end of the test, when the compound was becoming less effective, these waves were increasing in frequency and severity and were associated with increasing cough and nasal obstruction.

When the patient initially took capsules containing curcumin and black pepper, he experienced some gastro-esophageal discomfort and heartburn (VAS 2-3/10 for pain). However, when red pepper was associated with paracetamol heartburn increased (VAS 6) and required lansoprazole 30 mg. This discomfort lasted for around one to 2 hours.

Discussion

1- Strengths and limitations

N-of-1 trials appear to be well suited to find the rapid-onset treatments of a patient with COVID-19 to inform individualized treatment selection⁹. There are methodologically valid^{10 11}. In the present study, the patient had a stable condition assessed by repeated challenges with broccoli seeds and a rapid

washout ¹². The cough challenge is highly reproducible as shown for broccoli (10 open labelled in the pre-study, 4 DB-PC and 9 open labelled in the study). Moreover, it is very easy to count cough episodes. Nasal VAS may be less easy to assess but the patient has a much expertise on it.

One single patient was used for the challenges which has both strengths and weaknesses.

The choice of the nutrients was not easy but we tried to find Nrf2, TRPA1 and TRPV1 agonists as clear cut as possible. One important TRPA1 agonist is allicin that was not used since the patient is intolerant to garlic. Mustard oil alone cannot be found in a nutrient in France. Only a few challenges were carried out with the same nutrient but we wanted to study a larger number. The quality of the products raises other questions. The broccoli capsule was studied.

We do not have the exact composition of each nutrient and we used the labelled one.

The cough challenge does not reflect real life since inducing cough activates TRPA1/TRPV1 receptors that are sensitive to mechanical stress in a way that is not encountered in real life.

2- Optimal combination to be further tested in clinical trials

The results of this study are indicative of the efficacy of Nrf2 and TRPA1/TRPV1 agonists, but the best combination may not have been tested. The exact composition of each nutrient is required and its overall oxidative potency needs assessment. Then, different mixtures can be tested in controlled clinical trials. However, the results of the present study indicate that Nrf2 agonists or TRPA1/TRPV1 agonists alone are insufficient in the control of cough and nasal symptoms following COVID-19 infection. The combination was more effective and lasted for a longer period of time.

3- Possible putative mechanisms to be confirmed (Table 3)

Table 3: Interpretation of the results

	Effect observed	Tentative interpretation
1	All effective nutrients (except resveratrol and paracetamol) acting on induced cough and nasal symptoms are effective between 1 to 10 minutes.	This suggests a neurologic mechanism. In all conditions associated with COVID-19, there is an impaired redox homeostasis responsible for reactive oxygen species accumulation ¹ . TRPA1 is highly sensitive to oxidative stress ⁵² . Several COVID-19 symptoms may be induced by TRPA1/TRPV1 activation ^{7,53} . TRPA1/TRPV1 may be activated during COVID-19 and can be down-regulated very rapidly, possibly explaining the extremely swift onset of nutrients. Paracetamol was effective after 20 minutes since only its metabolites are TRPA1/TRPV1 agonists. ^{26,27}
2	Broccoli capsules that are effective on some symptoms of COVID-19 reduce nasal symptoms and induced cough from 5-10 minutes to around 4-6 hours.	Broccoli contains <ul style="list-style-type: none"> • relatively low levels of TRPA1/TRPV1 explaining its moderate effects by comparison to other nutrients for the first hour; • glucoraphanin and myrosinase that can produce sulforaphane ⁵ within 30 minutes ⁵⁴, the most potent Nrf2 natural agonist ⁵⁵, possibly explaining the slightly delayed and long-term effects.

3	Berberine, an Nrf2 agonist that does not have an apparent TRPA1 or TRPV1 interaction, is not effective on nasal symptoms and induced cough.	A moderate Nrf2 activity may be insufficient for the control of COVID-19 symptoms.
4	Zinc, at the recommended dose of 15 mg, is ineffective either as a single challenge (N=1) or in combination with broccoli and curcumin + green pepper (N=1)	The dose of Zinc may be too low.
5	Curcumin and black pepper reduce nasal symptoms and induced cough more rapidly, but their effect is not sustained after 2-3 hours.	Curcumin is a TRPA1 agonist ²¹ and black pepper a TRPA1 and TRV1 agonist ⁵⁶ (and both are Nrf2 agonists). Their potent and rapid effect is probably associated with the desensitization of the receptors.
6	There is a dose-dependent effect of curcumin and black pepper or red pepper. At the daily recommended dose, the capsules are more effective on nasal symptoms and induced cough than when using half a dose.	This is in line with receptor desensitization.
7	High doses of green tea reduce nasal symptoms and induced cough rapidly and for a short period (< 1 hr).	Polyphenols of green tea are TRPA1 agonists ^{22,47} (and Nrf2 agonists).
8	Resveratrol induces a delayed (30-45 min) reduction of nasal symptoms and induced cough.	Resveratrol is another TRPA1 agonist (and Nrf2 agonists) but its mechanisms of action may need some time to be effective ^{23,57} .
9	When broccoli (at half dose) is added to low-dose curcumin + black pepper, ginger or red pepper, high-dose green tea or regular dose resveratrol, the effects are very rapid and last for over 8 hrs.	The prolongation of the effects of both broccoli (sulforaphane) and TRAP1/TRPV1 agonists suggests an interaction between the two mechanisms. It is likely that TRAP1 desensitization is prolonged by the Nrf2 anti-oxidant effects of sulforaphane, leading to a down-regulation of the receptor.
10	The addition of paracetamol (low dose) to broccoli and curcumin + black pepper, red pepper prolongs the effects of the combination to over 14 hrs.	Some paracetamol metabolites (N-acetyl-p-benzoquinone imine) are TRPA1/TRPV1 agonists ²⁶ . During the cytokine storm in three patients, it was found that broccoli effects increased from around 6 to 10-12 hrs by paracetamol ¹⁴ . The prolongation of the effect reproduced in the current challenges is likely to be associated with TRAP1, since paracetamol takes some time to be metabolized.
11	Spontaneous cough is increased with the TRAP1/TRPV1 agonists by comparison to broccoli alone. These cough episodes (usually one or two mild coughs) occurred simultaneously with nasal obstruction and irritation of the larynx and trachea for less than one minute. They had no impact on the results of the following spontaneous challenge.	These episodes probably reflected the transient activation of the TRPA1/TRPV1 receptors followed by immediate desensitization.
12	At the end of the beneficial effect of the Nrf2 or TRPA1/TRPV1 -interacting nutrients, cough, nasal obstruction and tracheobronchial irritation increased very often simultaneously.	These findings suggest that TRPA1/TRPV1 receptors were active again.
13	At the end of the beneficial effect of the nutrients, a gastroesophageal discomfort occurred only with curcumin and black pepper. This effect lasted for 2-3 hrs and was very similar to symptoms occurring after wasabi ingestion when the patient ate sushi ⁵⁸ .	Wasabi is a potent TRPA1 agonist ⁵⁸ . These findings suggest the reactivation of TRPA1 receptors. There may be an interaction with TRVP1 receptors. The findings of 11-13 strongly suggest TRPA1/TRPV1 desensitization.

In the current study, the short duration of the effect of green tea or curcumin + black pepper or resveratrol suggests that TRP desensitization does not last long with the doses ingested. This is very important since, in Westernized diet, the amount of spices regularly consumed is usually low.

Desensitization of TRPV1 underlies the paradoxical analgesic effect of capsaicin. The TRPV1 receptors begin a refractory state, commonly termed as desensitization, that leads to the inhibition of receptor function⁵⁹. The acute desensitization of TRPV1 accounts for most of the reduction in responsiveness occurring within the first few (~20) seconds after the vanilloids are administered to the cell for the first time. Several signalling pathways including calcineurin, calmodulin or the decrease of Phosphatidylinositol 4,5-bisphosphate (PIP₂)⁶⁰ are involved in TRPV1 desensitization. TRPA1 is desensitized by homologous (mustard oil; a TRPA1 agonist) or heterologous (capsaicin; a TRPV1 agonist) agonists via Ca²⁺-independent and Ca²⁺-dependent pathways in the sensory neurons⁶¹. These rapid desensitization processes are likely to be involved in the rapid effects of TRPA1/V1 nutrients found in the current study.

Another form of TRP desensitization is 'tachyphylaxis', which is a reduction or the disappearance in the response to repeated applications of agonists^{62,63}. In Asian or Sub-Saharan countries, large amounts of spices are regularly eaten and it is likely that TRPA1/V1 receptors are permanently desensitized or have disappeared allowing people to eat large amounts. It is therefore possible that TRPA1/V1 tachyphylaxis may reduce the severity of COVID-19 symptoms, and even possibly reduce infection by SARS-CoV-2.

The increased duration of the effect when lower doses of curcumin + black pepper or red pepper were added to broccoli suggests that oxidative stress down-regulated by sulforaphane allows the desensitization of TRPs for a longer period of time. Moreover, for the first hour after ingestion, small waves of tracheal irritation, cough and nasal obstruction were perceived, suggesting activation-deactivation of the receptors. It probably takes one hour for sulforaphane to be synthesized and active in reducing the oxidative stress. This would indicate a cross talk between the anti-oxidative effect of Nrf2 and the desensitization of TRPA1/V1. This information is critical as it allows to propose low doses of TRPA1/V1 agonists that will be potentiated by Nrf2 agonists in COVID-19. The doses of agonists are still unclear and needs to be studied further.

4. Safety

The safety was clearly considered. The doses of broccoli were similar as those recommended as a food supplement (4 capsules a day). There is no safety issue with the doses administered. Curcumin and black pepper are routinely used in the Asian cuisine at relatively high doses and capsules are available in pharmacies or biofood shops. They appear to be safe although there was a warning issued by the ANSES (*agence nationale de sécurité sanitaire, alimentation, environnement, travail*, France) for curcumin as it may impact SARS-Cov-2 infection⁶⁴. However, ANSES recommends to use curcuma in the frame of a clinical trial. In 2010, the ANS (Food Additives and Nutrient Sources) of EFSA (European Food Safety Authority) re-evaluated the safety use of curcumin (E100) and established an acceptable daily intake (ADI) of 3 mg/kg of body weight based on the No-Observed-Adverse-Effect Level (NOAEL). This was re-evaluated in 2014⁶⁵. The same document stated that the dose of dietary food supplements supplied in a solid form including capsules should be 300 mg per day. All other nutrients were used at a dose far lower than the toxic dose.

Research should confirm these data and the intertwined mechanism of great importance in order to develop medications and foods based on TRPA1, TRPV1 and Nrf2 agonists. Moreover, more effective nutrients can be proposed.

References

1. Silvagno F, Vernone A, Pescarmona GP. The Role of Glutathione in Protecting against the Severe Inflammatory Response Triggered by COVID-19. *Antioxidants (Basel)*. 2020;9(7).
2. Martinez-Huelamo M, Rodriguez-Morato J, Boronat A, de la Torre R. Modulation of Nrf2 by Olive Oil and Wine Polyphenols and Neuroprotection. *Antioxidants (Basel)*. 2017;6(4).
3. Bousquet J, Czarlewski W, Blain H, Zuberbier T, Anto J. Rapid Response: Why Germany's case fatality rate seems so low: Is nutrition another possibility. *bmj*. 2020;<https://www.bmj.com/content/369/bmj.m1395/r-12>.
4. Bousquet J, Anto JM, Iaccarino G, et al. Is diet partly responsible for differences in COVID-19 death rates between and within countries? *Clin Transl Allergy*. 2020;10:16.
5. Bousquet J, Anto JM, Czarlewski W, et al. Cabbage and fermented vegetables: from death rate heterogeneity in countries to candidates for mitigation strategies of severe COVID-19. *Allergy*. 2020.
6. Talavera K, Startek JB, Alvarez-Collazo J, et al. Mammalian Transient Receptor Potential TRPA1 Channels: From Structure to Disease. *Physiol Rev*. 2020;100(2):725-803.
7. Bousquet J, Czarlewski W, Zuberbier T, et al. Spices to Control COVID-19 Symptoms: Yes, but Not Only. *Int Arch Allergy Immunol*. 2020:1-7.
8. Bousquet J, Czarlewski W, Zuberbier T, et al. Spices to control COVID-19: Yes, but not only.... *Int Arch Allergy Immunol*. 2020:in press.
9. Samuel J, Holder T, Molony D. N-of-1 Trials as a Decision Support Tool in Clinical Practice: A Protocol for a Systematic Literature Review and Narrative Synthesis. *Healthcare (Basel)*. 2019;7(4).
10. OCEBM Levels of Evidence - CEBM. CEBM. <https://www.cebmnet/2016/05/ocebm-levels-of-evidence/>. 2019.
11. Mitchell G, Nikles J. *The Essential Guide to N-of-1 Trials in Health*. Amsterdam: Springer Netherlands; 2015.
12. Bobe JR, De Freitas JK, Glicksberg BS. Exploring the Potential for Collaborative Use of an App-Based Platform for n-of-1 Trials Among Healthcare Professionals That Treat Patients With Insomnia. *Front Psychiatry*. 2020;11:530995.
13. Soldevila-Domenech N, Boronat A, Langohr K, de la Torre R. N-of-1 Clinical Trials in Nutritional Interventions Directed at Improving Cognitive Function. *Front Nutr*. 2019;6:110.
14. Bousquet J, Le-Moing V, Blain H, et al. Efficacy of broccoli and glucoraphanin in COVID-19: From hypothesis to proof-of-concept with three experimental clinical cases. *World Allergy Organ J*. [https://authors.elsevier.com/sd/article/S1939-4551\(20\)30401-4](https://authors.elsevier.com/sd/article/S1939-4551(20)30401-4). 2020.
15. Yagishita Y, Fahey JW, Dinkova-Kostova AT, Kensler TW. Broccoli or Sulforaphane: Is It the Source or Dose That Matters? *Molecules*. 2019;24(19).
16. Hindson J. Brassica vegetable metabolism by gut microbiota. *Nat Rev Gastroenterol Hepatol*. 2020;17(4):195.
17. Doheny-Adams T, Redeker K, Kittipol V, Bancroft I, SEHartley. Development of an efficient glucosinolate extraction method. *Plant methods*. 2017;13:17 DOI 10.1186/s13007-13017-10164-13008.
18. Bricker GV, Riedl KM, Ralston RA, Tober KL, Oberyshyn TM, Schwartz SJ. Isothiocyanate metabolism, distribution, and interconversion in mice following consumption of thermally processed broccoli sprouts or purified sulforaphane. *Mol Nutr Food Res*. 2014;58(10):1991-2000.
19. Vasanthi HR, Mukherjee S, Das DK. Potential health benefits of broccoli- a chemico-biological overview. *Mini Rev Med Chem*. 2009;9(6):749-759.
20. Fahey JW, Wade KL, Stephenson KK, et al. Bioavailability of Sulforaphane Following Ingestion of Glucoraphanin-Rich Broccoli Sprout and Seed Extracts with Active Myrosinase: A Pilot Study of the Effects of Proton Pump Inhibitor Administration. *Nutrients*. 2019;11(7).
21. Leamy AW, Shukla P, McAlexander MA, Carr MJ, Ghatta S. Curcumin ((E,E)-1,7-bis(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione) activates and desensitizes the nociceptor ion channel TRPA1. *Neurosci Lett*. 2011;503(3):157-162.
22. Kurogi M, Kawai Y, Nagatomo K, Tateyama M, Kubo Y, Saitoh O. Auto-oxidation products of epigallocatechin gallate activate TRPA1 and TRPV1 in sensory neurons. *Chem Senses*. 2015;40(1):27-46.
23. Nalli M, Ortar G, Moriello AS, Morera E, Di Marzo V, De Petrocellis L. TRPA1 channels as targets for resveratrol and related stilbenoids. *Bioorg Med Chem Lett*. 2016;26(3):899-902.
24. Huang TC, Chang WT, Hu YC, et al. Zinc Protects Articular Chondrocytes through Changes in Nrf2-Mediated Antioxidants, Cytokines and Matrix Metalloproteinases. *Nutrients*. 2018;10(4).
25. Hu H, Bandell M, Petrus MJ, Zhu MX, Patapoutian A. Zinc activates damage-sensing TRPA1 ion channels. *Nat Chem Biol*. 2009;5(3):183-190.
26. Andersson DA, Gentry C, Moss S, Bevan S. Transient receptor potential A1 is a sensory receptor for multiple products of oxidative stress. *J Neurosci*. 2008;28(10):2485-2494.
27. Wang Y, Lin W, Wu N, et al. An insight into paracetamol and its metabolites using molecular docking and molecular dynamics simulation. *J Mol Model*. 2018;24(9):243.
28. Copple IM, Goldring CE, Jenkins RE, et al. The hepatotoxic metabolite of acetaminophen directly activates the Keap1-Nrf2 cell defense system. *Hepatology*. 2008;48(4):1292-1301.
29. Ibarra Y, Blair NT. Benzoquinone reveals a cysteine-dependent desensitization mechanism of TRPA1. *Mol Pharmacol*. 2013;83(5):1120-1132.
30. Wojdyla K, Wrzesinski K, Williamson J, Fey SJ, Rogowska-Wrzesinska A. Acetaminophen-induced S-nitrosylation and S-sulfonylation signalling in 3D cultured hepatocarcinoma cell spheroids. *Toxicol Res (Camb)*. 2016;5(3):905-920.

31. Bousquet PJ, Combescure C, Neukirch F, et al. Visual analog scales can assess the severity of rhinitis graded according to ARIA guidelines. *Allergy*. 2007;62(4):367-372.
32. Klimek L, Bergmann KC, Biedermann T, et al. Visual analogue scales (VAS): Measuring instruments for the documentation of symptoms and therapy monitoring in cases of allergic rhinitis in everyday health care: Position Paper of the German Society of Allergology (AeDA) and the German Society of Allergy and Clinical Immunology (DGAKI), ENT Section, in collaboration with the working group on Clinical Immunology, Allergology and Environmental Medicine of the German Society of Otorhinolaryngology, Head and Neck Surgery (DGHNOKHC). *Allergo J Int*. 2017;26(1):16-24.
33. Spinou A, Birring SS. An update on measurement and monitoring of cough: what are the important study endpoints? *J Thorac Dis*. 2014;6(Suppl 7):S728-734.
34. Malavolta M, Bracci M, Santarelli L, et al. Inducers of Senescence, Toxic Compounds, and Senolytics: The Multiple Faces of Nrf2-Activating Phytochemicals in Cancer Adjuvant Therapy. *Mediators Inflamm*. 2018;2018:4159013.
35. Saw CL, Guo Y, Yang AY, et al. The berry constituents quercetin, kaempferol, and pterostilbene synergistically attenuate reactive oxygen species: involvement of the Nrf2-ARE signaling pathway. *Food Chem Toxicol*. 2014;72:303-311.
36. Nakamura T, Miyoshi N, Ishii T, Nishikawa M, Ikushiro S, Watanabe T. Activation of transient receptor potential ankyrin 1 by quercetin and its analogs. *Biosci Biotechnol Biochem*. 2016;80(5):949-954.
37. Ashrafizadeh M, Fekri HS, Ahmadi Z, Farkhondeh T, Samarghandian S. Therapeutic and biological activities of berberine: The involvement of Nrf2 signaling pathway. *J Cell Biochem*. 2020;121(2):1575-1585.
38. Zan Y, Kuai CX, Qiu ZX, Huang F. Berberine Ameliorates Diabetic Neuropathy: TRPV1 Modulation by PKC Pathway. *Am J Chin Med*. 2017;45(8):1709-1723.
39. Patel SS, Acharya A, Ray RS, Agrawal R, Raghuvanshi R, Jain P. Cellular and molecular mechanisms of curcumin in prevention and treatment of disease. *Crit Rev Food Sci Nutr*. 2020;60(6):887-939.
40. Nalli M, Ortar G, Schiano Moriello A, Di Marzo V, De Petrocellis L. Effects of curcumin and curcumin analogues on TRP channels. *Fitoterapia*. 2017;122:126-131.
41. Joung EJ, Li MH, Lee HG, et al. Capsaicin induces heme oxygenase-1 expression in HepG2 cells via activation of PI3K-Nrf2 signaling: NAD(P)H:quinone oxidoreductase as a potential target. *Antioxid Redox Signal*. 2007;9(12):2087-2098.
42. Moran MM, Szallasi A. Targeting nociceptive transient receptor potential channels to treat chronic pain: current state of the field. *Br J Pharmacol*. 2018;175(12):2185-2203.
43. Yang F, Zheng J. Understand spiciness: mechanism of TRPV1 channel activation by capsaicin. *Protein Cell*. 2017;8(3):169-177.
44. de Lima RMT, Dos Reis AC, de Menezes APM, et al. Protective and therapeutic potential of ginger (*Zingiber officinale*) extract and [6]-gingerol in cancer: A comprehensive review. *Phytother Res*. 2018;32(10):1885-1907.
45. Yang MQ, Ye LL, Liu XL, et al. Gingerol activates noxious cold ion channel TRPA1 in gastrointestinal tract. *Chin J Nat Med*. 2016;14(6):434-440.
46. Yin Y, Dong Y, Vu S, et al. Structural mechanisms underlying activation of TRPV1 channels by pungent compounds in gingers. *Br J Pharmacol*. 2019;176(17):3364-3377.
47. Peixoto-Neves D, Soni H, Adebisi A. CGRPergic Nerve TRPA1 Channels Contribute to Epigallocatechin Gallate-Induced Neurogenic Vasodilation. *ACS Chem Neurosci*. 2019;10(1):216-220.
48. Li Z, Zhang J, Ren X, Liu Q, Yang X. The mechanism of quercetin in regulating osteoclast activation and the PAR2/TRPV1 signaling pathway in the treatment of bone cancer pain. *Int J Clin Exp Pathol*. 2018;11(11):5149-5156.
49. Srinivasan K. Biological Activities of Red Pepper (*Capsicum annuum*) and Its Pungent Principle Capsaicin: A Review. *Crit Rev Food Sci Nutr*. 2016;56(9):1488-1500.
50. Yu L, Wang S, Kogure Y, Yamamoto S, Noguchi K, Dai Y. Modulation of TRP channels by resveratrol and other stilbenoids. *Mol Pain*. 2013;9:3.
51. Luo J, Bavencoffe A, Yang P, et al. Zinc Inhibits TRPV1 to Alleviate Chemotherapy-Induced Neuropathic Pain. *J Neurosci*. 2018;38(2):474-483.
52. Kadkova A, Synytsya V, Krusek J, Zimova L, Vlachova V. Molecular basis of TRPA1 regulation in nociceptive neurons. A review. *Physiol Res*. 2017;66(3):425-439.
53. Bousquet J, Cristol JP, Czarlewski W, et al. Nrf2-interacting nutrients and COVID-19: time for research to develop adaptation strategies. *Clin Transl Allergy*. 2020;10(1):58.
54. Gasper AV, Al-Janobi A, Smith JA, et al. Glutathione S-transferase M1 polymorphism and metabolism of sulforaphane from standard and high-glucosinolate broccoli. *Am J Clin Nutr*. 2005;82(6):1283-1291.
55. Silva-Islas CA, Maldonado PD. Canonical and non-canonical mechanisms of Nrf2 activation. *Pharmacol Res*. 2018;134:92-99.
56. Okumura Y, Narukawa M, Iwasaki Y, et al. Activation of TRPV1 and TRPA1 by black pepper components. *Biosci Biotechnol Biochem*. 2010;74(5):1068-1072.
57. Nakao S, Mabuchi M, Wang S, et al. Synthesis of resveratrol derivatives as new analgesic drugs through desensitization of the TRPA1 receptor. *Bioorg Med Chem Lett*. 2017;27(14):3167-3172.
58. Terada Y, Masuda H, Watanabe T. Structure-Activity Relationship Study on Isothiocyanates: Comparison of TRPA1-Activating Ability between Allyl Isothiocyanate and Specific Flavor Components of Wasabi, Horseradish, and White Mustard. *J Nat Prod*. 2015;78(8):1937-1941.

59. Nilius B, Appendino G. Spices: the savory and beneficial science of pungency. *Rev Physiol Biochem Pharmacol.* 2013;164:1-76.
60. Rohacs T. Phosphoinositide regulation of TRPV1 revisited. *Pflugers Arch.* 2015;467(9):1851-1869.
61. Akopian AN, Ruparel NB, Jeske NA, Hargreaves KM. Transient receptor potential TRPA1 channel desensitization in sensory neurons is agonist dependent and regulated by TRPV1-directed internalization. *J Physiol.* 2007;583(Pt 1):175-193.
62. Touska F, Marsakova L, Teisinger J, Vlachova V. A "cute" desensitization of TRPV1. *Curr Pharm Biotechnol.* 2011;12(1):122-129.
63. Sanz-Salvador L, Andres-Borderia A, Ferrer-Montiel A, Planells-Cases R. Agonist- and Ca²⁺-dependent desensitization of TRPV1 channel targets the receptor to lysosomes for degradation. *J Biol Chem.* 2012;287(23):19462-19471.
64. Risques liés à la consommation de compléments alimentaires contenant des plantes pouvant interférer avec la réponse immunitaire et inflammatoire associée à l'infection par le SARS-CoV-2. *anses, sisine* n°2020-SA-0045. 2020;<https://www.anses.fr/fr/system/files/NUT2020SA0045.pdf>.
65. Refined exposure assessment for curcumin (E 100). European Food Safety Authority. *EFSA Journal.* 2014;12(10):<https://efsa.onlinelibrary.wiley.com/doi/pdf/10.2903/j.efsa.2014.3876>.